

Simulations of Weighted Tree Automata

Zoltán Ésik^{1,*} and Andreas Maletti^{2,**}

¹ University of Szeged, Department of Computer Science
Árpád tér 2, 6720 Szeged, Hungary. ze@inf.u-szeged.hu

² Universitat Rovira i Virgili, Departament de Filologies Romàniques
Avinguda de Catalunya 35, 43002 Tarragona, Spain. andreas.maletti@urv.cat

Abstract. Simulations of weighted tree automata (wta) are considered. It is shown how such simulations can be decomposed into simpler functional and dual functional simulations also called forward and backward simulations. In addition, it is shown in several cases (fields, commutative rings, NOETHERIAN semirings, semiring of natural numbers) that all equivalent wta M and N can be joined by a finite chain of simulations. More precisely, in all mentioned cases there is a single wta that simulates both M and N . Those results immediately yield decidability of equivalence provided that the semiring is finitely (and effectively) presented.

1 Introduction

Weighted tree automata are widely used in applications such as model checking [1] and natural language processing [21]. They finitely represent mappings, called tree series, that assign a weight to each tree. For example, a probabilistic parser would return a tree series that assigns to each parse tree its likelihood. Consequently, several toolkits [20, 24, 10] implement weighted tree automata.

The notion of simulation that is used in this paper is a generalization of the simulations for unweighted and weighted (finite) string automata of [6, 15]. The aim is to relate structurally equivalent automata. The results of [6, Section 9.7] and [22] show that two unweighted string automata (i.e., potentially nondeterministic string automata over the BOOLEAN semiring) are equivalent if and only if they can be connected by a finite chain of relational simulations, and that in fact *functional* and *dual functional* simulations are sufficient. Simulations for weighted string automata (wsa) are called *conjugacies* in [3, 4], where it is shown that for all fields, many rings including the ring \mathbb{Z} of integers, and the semiring \mathbb{N} of natural numbers, two wsa are equivalent if and only if they can be connected by a finite chain of simulations. It is also shown that even a finite

* Partially supported by grant no. K 75249 from the *National Foundation of Hungary for Scientific Research* and by the TÁMOP-4.2.2/08/1/2008-0008 program of the *Hungarian National Development Agency*.

** Financially supported by the *Ministerio de Educación y Ciencia* grants JDCI-2007-760 and MTM-2007-63422 and the *European Science Foundation* short-visit grant 2978 in the activity “Automata: from Mathematics to Applications”.

chain of functional (*covering*) and dual functional (*co-covering*) simulations is sufficient. The origin of those results can be traced back to the pioneering work of SCHÜTZENBERGER in the early 60's, who proved that every wsa over a field is equivalent to a minimal wsa that is simulated by every *trim* equivalent wsa [5]. Relational simulations of wsa are also studied in [9], where they are used to reduce the size of wsa. The relationship between functional simulations and the MILNER-PARK notion of bisimulation [25, 26] is discussed in [6, 9].

In this contribution, we investigate simulations for weighted (finite) tree automata (wta). SCHÜTZENBERGER's minimization method was extended to wta over fields in [2, 8]. In addition, relational and functional simulations for wta are probably first used in [12, 13, 18]. Moreover, simulations can be generalized to presentations in algebraic theories [6], which seems to cover all mentioned instances. Here, we extend and improve the results of [3, 4] to wta. In particular, we show that two wta over a commutative ring, NOETHERIAN semiring, or the semiring \mathbb{N} are equivalent if and only if they are connected by a finite chain of simulations. Moreover, we discuss when the simulations can be replaced by functional and dual functional simulations, which are efficiently computable [18]. Such results are important because they immediately yield the decidability of equivalence provided that the semiring is finitely and effectively presented.

2 Preliminaries

The set of nonnegative integers is \mathbb{N} . For every $k \in \mathbb{N}$, the set $\{i \in \mathbb{N} \mid 1 \leq i \leq k\}$ is simply denoted by $[k]$. We write $|A|$ for the cardinality of the set A . A *semiring* is an algebraic structure $\mathcal{A} = (A, +, \cdot, 0, 1)$ such that $(A, +, 0)$ and $(A, \cdot, 1)$ are monoids, of which the former is commutative, and \cdot distributes both-sided over finite sums (i.e., $a \cdot 0 = 0 = 0 \cdot a$ for every $a \in A$ and $a \cdot (b + c) = ab + ac$ and $(b + c) \cdot a = ba + ca$ for every $a, b, c \in A$). The semiring \mathcal{A} is *commutative* if $(A, \cdot, 1)$ is commutative. It is a *ring* if there exists an element $-1 \in A$ such that $1 + (-1) = 0$. The set U is the set $\{a \in A \mid \exists b \in A: ab = 1 = ba\}$ of (*multiplicative*) *units*. The semiring \mathcal{A} is a *semifield* if $U = A \setminus \{0\}$; i.e., for every $a \in A$ there exists a *multiplicative inverse* $a^{-1} \in A$ such that $aa^{-1} = 1 = a^{-1}a$. A *field* is a semifield that is also a ring. Let $\langle B \rangle_+ = \{b_1 + \dots + b_n \mid n \in \mathbb{N}, b_1, \dots, b_n \in B\}$ for every $B \subseteq A$. If $A = \langle B \rangle_+$, then \mathcal{A} is *additively generated by B*. The semiring \mathcal{A} is *equisubtractive* if for every $a_1, a_2, b_1, b_2 \in A$ with $a_1 + b_1 = a_2 + b_2$ there exist $c_1, c_2, d_1, d_2 \in A$ such that (i) $a_1 = c_1 + d_1$, (ii) $b_1 = c_2 + d_2$, (iii) $a_2 = c_1 + c_2$, and (iv) $b_2 = d_1 + d_2$. It is *zero-sum free* (*zero-divisor free*, respectively) if $a + b = 0$ ($a \cdot b = 0$, respectively) implies $0 \in \{a, b\}$ for every $a, b \in A$. Finally, it is *positive* if it is both zero-sum and zero-divisor free. Clearly, any nontrivial (i.e., $0 \neq 1$) ring is not zero-sum free, and any semifield is zero-divisor free. An infinitary sum operation \sum is a family $(\sum_I)_I$ such that $\sum_I: A^I \rightarrow A$. We generally write $\sum_{i \in I} a_i$ instead of $\sum_I (a_i)_{i \in I}$. The semiring \mathcal{A} together with the infinitary sum operation \sum is *complete* [11, 17, 19] if for all index sets I and $(a_i)_{i \in I} \in A^I$

- $\sum_{i \in I} a_i = a_{j_1} + a_{j_2}$ if $I = \{j_1, j_2\}$ with $j_1 \neq j_2$,
- $\sum_{i \in I} a_i = \sum_{j \in J} (\sum_{i \in I_j} a_i)$ for every partition $(I_j)_{j \in J}$ of I , and

$$- a \cdot (\sum_{i \in I} a_i) = \sum_{i \in I} aa_i \text{ and } (\sum_{i \in I} a_i) \cdot a = \sum_{i \in I} a_i a \text{ for every } a \in A.$$

An \mathcal{A} -semimodule is a commutative monoid $(B, +, 0)$ together with an action $\cdot : A \times B \rightarrow B$, written as juxtaposition, such that for every $a, a' \in A$ and $b, b' \in B$ we have (i) $(a + a')b = ab + a'b$, (ii) $a(b + b') = ab + ab'$, (iii) $0b = 0$, (iv) $1b = b$, and (v) $(a \cdot a')b = a(a'b)$. The semiring \mathcal{A} is NOETHERIAN if all subsemimodules of every finitely-generated \mathcal{A} -semimodule are again finitely-generated.

In the following, we identify index sets of equal cardinality. Let $X \in A^{I_1 \times J_1}$ and $Y \in A^{I_2 \times J_2}$ for finite sets I_1, I_2, J_1, J_2 . We use upper-case letters (like C, D, E, X, Y) for matrices and the corresponding lower-case letters for their entries. The matrix X is *relational* if $X \in \{0, 1\}^{I_1 \times J_1}$. Clearly, a relational X corresponds to a relation $\rho_X \subseteq I_1 \times J_1$ (and vice versa) by $(i, j) \in \rho_X$ if and only if $x_{ij} = 1$. Moreover, a relational matrix X is *functional*, *surjective*, or *injective* if ρ_X has this property. As usual, we denote the *transpose* of X by X^T , and we call X *nondegenerate* if it has no rows or columns of entirely zeroes. A *diagonal* matrix X is such that $x_{ij} = 0$ for every $i \neq j$. Finally, the matrix X is invertible if there exists a matrix X^{-1} such that $XX^{-1} = I = X^{-1}X$ where I is the unit matrix. The KRONECKER product $X \otimes Y \in A^{(I_1 \times I_2) \times (J_1 \times J_2)}$ is such that $(X \otimes Y)_{i_1 i_2, j_1 j_2} = x_{i_1 j_1} y_{i_2 j_2}$ for every $i_1 \in I_1, i_2 \in I_2, j_1 \in J_1$, and $j_2 \in J_2$. We let $X^{0, \otimes} = (1)$ and $X^{i+1, \otimes} = X^{i, \otimes} \otimes X$ for every $i \in \mathbb{N}$.

Finally, let us move to trees. A *ranked alphabet* is a finite set Σ together with a mapping $\text{rk} : \Sigma \rightarrow \mathbb{N}$. We often just write Σ for a ranked alphabet and assume that the mapping rk is implicit. We write $\Sigma_k = \{\sigma \in \Sigma \mid \text{rk}(\sigma) = k\}$ for the set of all k -ary symbols. The set of Σ -trees is the smallest set T_Σ such that $\sigma(t_1, \dots, t_k) \in T_\Sigma$ for all $\sigma \in \Sigma_k$ and $t_1, \dots, t_k \in T_\Sigma$. A *tree series* is a mapping $\varphi : T_\Sigma \rightarrow A$. The set of all such tree series is denoted by $A\langle\langle T_\Sigma \rangle\rangle$. For every $\varphi \in A\langle\langle T_\Sigma \rangle\rangle$ and $t \in T_\Sigma$, we often write (φ, t) instead of $\varphi(t)$.

A *weighted tree automaton (over \mathcal{A})*, for short: wta, is a system (Σ, Q, μ, F) with an input ranked alphabet Σ , a finite set Q of *states*, transitions $\mu = (\mu_k)_{k \in \mathbb{N}}$ such that $\mu_k : \Sigma_k \rightarrow A^{Q^k \times Q}$ for every $k \in \mathbb{N}$, and a *final weight vector* $F \in A^Q$. Next, let us introduce the semantics $\|M\|$ of the wta M . We first define the mapping $h_\mu : T_\Sigma \rightarrow A^Q$ for every $\sigma \in \Sigma_k$ and $t_1, \dots, t_k \in T_\Sigma$ by $h_\mu(\sigma(t_1, \dots, t_k)) = (h_\mu(t_1) \otimes \dots \otimes h_\mu(t_k)) \cdot \mu_k(\sigma)$, where the final product \cdot is the classical matrix product. Then $(\|M\|, t) = h_\mu(t)F$ for every $t \in T_\Sigma$, where the product is the usual inner (dot) product. The wta M is *trim* if every state is accessible and co-accessible in the BOOLEAN wta obtained by replacing every nonzero weight by 1.




Fig. 1. Illustration of simulation.

3 Simulation

Let us introduce the main notion of the paper. From now on, let $M = (\Sigma, Q, \mu, F)$ and $N = (\Sigma, P, \nu, G)$ be wta. Then M *simulates* N (cf., [6, 15], [3, Def. 1], and

[12, Def. 35]) if there is $X \in A^{Q \times P}$ such that $F = XG$ and $\mu_k(\sigma)X = X^{k, \otimes} \cdot \nu_k(\sigma)$ for every $\sigma \in \Sigma_k$. The matrix X is called *transfer matrix*, and we write $M \rightarrow_X N$ if M simulates N with transfer matrix X . Note that $X_{i_1 \dots i_k, j_1 \dots j_k}^{k, \otimes} = \prod_{\ell=1}^k x_{i_\ell, j_\ell}$. We illustrate the notion of simulation in Fig. 1. If $M \rightarrow_X M'$ and $M' \rightarrow_Y N$, then $M \rightarrow_{XY} N$. Moreover, if $M \rightarrow_X N$, then M and N are equivalent.

Theorem 1. *If M simulates N , then M and N are equivalent.*

Next, we prepare the result for functional simulations. To this end, we first need to prove in which cases the transfer matrix is nondegenerate.

Lemma 2. *Let M and N be trim and $M \rightarrow_X N$. If (i) X is functional or (ii) \mathcal{A} is positive, then X is nondegenerate.*

Now we relate functional simulation to forward simulation [18, Def. 1]. A surjective mapping $\rho: Q \rightarrow P$ is a *forward simulation* from M to N if (i) $F_q = G_{\rho(q)}$ for every $q \in Q$ and (ii) $\sum_{q \in \rho^{-1}(p)} \mu_k(\sigma)_{q_1 \dots q_k, q} = \nu_k(\sigma)_{\rho(q_1) \dots \rho(q_k), p}$ for every $p \in P$, $\sigma \in \Sigma_k$, and $q_1, \dots, q_k \in Q$. We say that M *forward simulates* N , written $M \rightarrow N$, if there exists a forward simulation from M to N . Similarly, we can relate backward simulation [18, Def. 16] to dual functional simulation. A surjective function $\rho: Q \rightarrow P$ is a *backward simulation* from M to N if $\sum_{q \in \rho^{-1}(p)} F_q = G_p$ for every $p \in P$ and $\sum_{q_1 \in \rho^{-1}(p_1), \dots, q_k \in \rho^{-1}(p_k)} \mu_k(\sigma)_{q_1 \dots q_k, q} = \nu_k(\sigma)_{p_1 \dots p_k, \rho(q)}$ for every $q \in Q$, $\sigma \in \Sigma_k$, and $p_1, \dots, p_k \in P$. We say that M *backward simulates* N , written $M \leftarrow N$, if there exists a backward simulation from M to N . Using Lemma 2 we obtain the following statement.

Lemma 3. *Let N be trim. Then $M \rightarrow N$ if and only if there exists a functional transfer matrix X such that $M \rightarrow_X N$. Moreover, $M \leftarrow N$ if and only if there exists a transfer matrix X such that X^T is functional and $N \rightarrow_X M$.*

Next, we recall two important matrix decomposition results of [3].

Lemma 4. *If $A = \langle U \rangle_+$, then for every $X \in A^{Q \times P}$ there exist matrices C, E, D such that (i) $X = CED$, (ii) C^T and D are functional, and (iii) E is an invertible diagonal matrix. If (a) X is nondegenerate or (b) \mathcal{A} has (nontrivial) zero-sums, then C^T and D can be chosen to be surjective.*

Lemma 5. *Let \mathcal{A} be equisubtractive. Moreover, let $R \in A^Q$ and $C \in A^P$ be such that $\sum_{q \in Q} r_q = \sum_{p \in P} c_p$. Then there exists a matrix $X \in A^{Q \times P}$ with row sums R and column sums C .*

Using all the previous results, we can now obtain the main result of this section, which shows how we can decompose simulation into functional and dual functional simulation (or: forward and backward simulation, respectively).

Theorem 6. *Let \mathcal{A} be equisubtractive with $A = \langle U \rangle_+$. Then $M \rightarrow_X N$ if and only if there exist two wta M' and N' such that (i) $M \rightarrow_C M'$ where C^T is functional, (ii) $M' \rightarrow_E N'$ where E is an invertible diagonal matrix, and (iii) $N' \rightarrow_D N$ where D is functional. If M and N are trim, then $M' \leftarrow M$ and $N' \rightarrow N$.*

Proof. Clearly, $M \rightarrow_C M' \rightarrow_E N' \rightarrow_D N$, which proves that $M \rightarrow_{CED} N$. For the converse, Lemma 4 shows that there exist matrices C , E , and D such that $X = CED$, C^T and D are functional matrices, and $E \in A^{I \times I}$ is an invertible diagonal matrix. Finally, let $\varphi: I \rightarrow Q$ and $\psi: I \rightarrow P$ be the functions associated to C^T and D .

It remains to determine the wta M' and N' . Let $M' = (\Sigma, I, \mu', F')$ and $N' = (\Sigma, I, \nu', G')$ with $G' = DG$ and $F' = EDG$. Then we have $CF' = CEDG = XG = F$. Thus, it remains to specify $\mu'_k(\sigma)$ and $\nu'_k(\sigma)$ for every $\sigma \in \Sigma_k$. To this end, we determine a matrix $Y \in A^{I^k \times I}$ such that we have (1) $C^{k, \otimes} \cdot Y = \mu_k(\sigma)CE$ and (2) $YD = E^{k, \otimes} \cdot D^{k, \otimes} \cdot \nu_k(\sigma)$. Let $\mu'_k(\sigma) = YE^{-1}$ and $\nu'_k(\sigma) = (E^{k, \otimes})^{-1} \cdot Y$. Consequently, we have $\mu_k(\sigma)C = C^{k, \otimes} \cdot \mu'_k(\sigma)$, $\mu'_k(\sigma)E = E^{k, \otimes} \cdot \nu'_k(\sigma)$, and $\nu'_k(\sigma)D = D^{k, \otimes} \cdot \nu_k(\sigma)$. These equalities are displayed in Fig. 2 (right).

Finally, we need to specify the matrix Y . For every $q \in Q$ and $p \in P$, let $I_q = \varphi^{-1}(q)$ and $J_p = \psi^{-1}(p)$. Obviously, Y can be decomposed into disjoint (not necessarily contiguous) submatrices $Y_{q_1 \dots q_k, p} \in A^{(I_{q_1} \times \dots \times I_{q_k}) \times J_p}$ with $q_1, \dots, q_k \in Q$ and $p \in P$. Then properties (1) and (2) hold if and only if for every $q_1, \dots, q_k \in Q$ and $p \in P$ the following two conditions hold:

1. For every $i \in I$ such that $\psi(i) = p$, the sum of the i -column of $Y_{q_1 \dots q_k, p}$ is $\mu_k(\sigma)_{q_1 \dots q_k, \varphi(i)} \cdot e_{i, i}$.
2. For all $i_1, \dots, i_k \in I$ such that $\varphi(i_j) = q_j$ for every $j \in [k]$, the sum of the (i_1, \dots, i_k) -row of $Y_{q_1 \dots q_k, p}$ is $\prod_{j=1}^k e_{i_j, i_j} \cdot \nu_k(\sigma)_{\psi(i_1) \dots \psi(i_k), p}$.

Those two conditions are compatible because

$$\begin{aligned} & \sum_{\substack{i \in I \\ \psi(i)=p}} \mu_k(\sigma)_{q_1 \dots q_k, \varphi(i)} \cdot e_{i, i} = (\mu_k(\sigma)CED)_{q_1 \dots q_k, p} = (\mu_k(\sigma)X)_{q_1 \dots q_k, p} \\ & = (X^{k, \otimes} \cdot \nu_k(\sigma))_{q_1 \dots q_k, p} = (C^{k, \otimes} \cdot E^{k, \otimes} \cdot D^{k, \otimes} \cdot \nu_k(\sigma))_{q_1 \dots q_k, p} \\ & = \sum_{\substack{i_1, \dots, i_k \in I \\ \forall j \in [k]: \varphi(i_j)=q_j}} \left(\prod_{j=1}^k e_{i_j, i_j} \right) \cdot \nu_k(\sigma)_{\psi(i_1) \dots \psi(i_k), p} . \end{aligned}$$

Consequently, the row and column sums of the submatrices $Y_{q_1 \dots q_k, p}$ are consistent, which yields that we can determine all the submatrices (and thus the whole matrix) by Lemma 5. If M and N are trim, then either (a) \mathcal{A} is zero-sum free (and thus positive because it is additively generated by its units), in which case X is nondegenerate by Lemma 2, or (b) \mathcal{A} has nontrivial zero-sums. In both cases, Lemma 4 shows that the matrices C^T and D are surjective, which yields the additional statement by Lemma 3. \square




Fig. 2. Relations between the matrices.

The decomposition of simulations into forward and backward simulation is effective and has computational benefits because it is shown in [18] that forward and backward simulation can be efficiently computed. To keep the presentation simple, we will continue to deal with simulation in the following. However, in many of the following cases they can be decomposed.

4 Category of simulations

In this section our aim is to show that several well-known constructions of wta are *functorial*: they may be extended to simulations in a functorial way. Below we will only deal with the sum, HADAMARD product, σ_0 -product, and σ_0 -iteration (cf. [14]). Scalar OI-substitution, \dagger (the dagger operation) [7], homomorphism, quotient, and top-concatenation [14] may be covered in a similar fashion.

In this section, let \mathcal{A} be commutative. Moreover, let $M = (\Sigma, Q, \mu, F)$, $M' = (\Sigma, Q', \mu', F')$, and $M'' = (\Sigma, Q'', \mu'', F'')$ be wta. We already remarked that, if $M \rightarrow_X M'$ and $M' \rightarrow_Y M''$, then $M \rightarrow_{XY} M''$. Moreover, $M \rightarrow_I M$ with the unit matrix $I \in A^{Q \times Q}$. Thus, wta over the alphabet Σ form a category \mathbf{Sim}_Σ .

In the following, let $M = (\Sigma, Q, \mu, F)$ and $N = (\Sigma, P, \nu, G)$ be wta such that $Q \cap P = \emptyset$. The sum $M + N$ of M and N is the wta $(\Sigma, Q \cup P, \kappa, H)$ where $H = \langle F, G \rangle$ and

$$\kappa_k(\sigma)_{q_1 \dots q_k, q} = (\mu_k(\sigma) + \nu_k(\sigma))_{q_1 \dots q_k, q} = \begin{cases} \mu_k(\sigma)_{q_1 \dots q_k, q} & \text{if } q, q_1, \dots, q_k \in Q \\ \nu_k(\sigma)_{q_1 \dots q_k, q} & \text{if } q, q_1, \dots, q_k \in P \\ 0 & \text{otherwise.} \end{cases}$$

for all $\sigma \in \Sigma_k$ and $q, q_1, \dots, q_k \in Q \cup P$. It is known that $\|M + N\| = \|M\| + \|N\|$. Next, we extend the sum construction to simulations. To this end, let $M \rightarrow_X M'$ and $N \rightarrow_Y N'$ with $N' = (\Sigma, P', \nu', G')$. The sum $X + Y \in A^{(Q \cup P) \times (Q' \cup P')}$ of the transfer matrices X and Y is $X + Y = \begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}$. Then $(M + N) \rightarrow_{X+Y} (M' + N')$.

Proposition 7. *The function $+$, which is defined on wta and transfer matrices, is a functor $\mathbf{Sim}_\Sigma^2 \rightarrow \mathbf{Sim}_\Sigma$.*

Next, we treat the remaining operations. Let σ_0 be a distinguished symbol in Σ_0 . The σ_0 -product $M \cdot_{\sigma_0} N$ of M with N is the wta $(\Sigma, Q \cup P, \kappa, H)$ such that $H = \langle F, 0 \rangle$ and for each $\sigma \in \Sigma_k$ with $\sigma \neq \sigma_0$,

$$\kappa_k(\sigma)_{q_1 \dots q_k, q} = \begin{cases} \mu_k(\sigma)_{q_1 \dots q_k, q} & \text{if } q, q_1, \dots, q_k \in Q \\ \mu_0(\sigma_0)_q \cdot \sum_{p \in P} \nu_k(\sigma)_{q_1 \dots q_k, p} G_p & \text{if } q \in Q \text{ and } q_1, \dots, q_k \in P \\ \nu_k(\sigma)_{q_1 \dots q_k, q} & \text{if } q, q_1, \dots, q_k \in P \\ 0 & \text{otherwise.} \end{cases}$$

Moreover,

$$\kappa_0(\sigma_0)_q = \begin{cases} \mu_0(\sigma_0)_q \cdot \sum_{p \in P} \nu_0(\sigma_0)_p G_p & \text{if } q \in Q \\ \nu_0(\sigma_0)_q & \text{if } q \in P. \end{cases}$$

It is known that $\|M \cdot_{\sigma_0} N\| = \|M\| \cdot_{\sigma_0} \|N\|$. Let $M \rightarrow_X M'$ and $N \rightarrow_Y N'$. We define $X \cdot_{\sigma_0} Y = X + Y$. The HADAMARD product $M \cdot_{\mathbb{H}} N$ is the wta $(\Sigma, Q \times P, \kappa, H)$ where $H = F \otimes G$ and $\kappa_k(\sigma) = \mu_k(\sigma) \otimes \nu_k(\sigma)$ for all $\sigma \in \Sigma_k$. If $M \rightarrow_X M'$ and $N \rightarrow_Y N'$, then we define $X \cdot_{\mathbb{H}} Y = X \otimes Y$. Finally, let \mathcal{A} be complete. Thus, \mathcal{A} allows the definition of the star operation $a^* = \sum_{n \in \mathbb{N}} a^n$ for every $a \in A$. The σ_0 -iteration $M^{*\sigma_0}$ of M is the wta (Σ, Q, κ, F) where

$$\kappa_k(\sigma)_{q_1 \dots q_k, q} = \mu_k(\sigma)_{q_1 \dots q_k, q} + \|M\|(\sigma_0)^* \cdot \sum_{p \in Q} \mu_k(\sigma)_{q_1 \dots q_k, p} F_p$$

for all $\sigma \in \Sigma_k \setminus \{\sigma_0\}$ and $\kappa_0(\sigma_0) = \mu_0(\sigma_0)$. If $M \rightarrow_X M'$, then we define $X^{*\sigma_0} = X$.

Proposition 8. *The functions \cdot_{σ_0} and $\cdot_{\mathbb{H}}$, which are defined on wta and transfer matrices, are functors $\mathbf{Sim}_{\Sigma}^2 \rightarrow \mathbf{Sim}_{\Sigma}$. Moreover, σ_0 -iteration is a functor $\mathbf{Sim}_{\Sigma} \rightarrow \mathbf{Sim}_{\Sigma}$ if \mathcal{A} is complete.*

5 Joint reduction

In this section, we will establish equivalence results using an improved version of the approach called *joint reduction* in [4]. Let $V \subseteq A^I$ be a set of vectors for a finite set I . The \mathcal{A} -semimodule generated by V is denoted by $\langle V \rangle$. Given two wta $M = (\Sigma, Q, \mu, F)$ and $N = (\Sigma, P, \nu, G)$ with $Q \cap P = \emptyset$, we first compute $M + N = (\Sigma, Q \cup P, \mu', F')$ as defined in Section 4. The aim is to compute a finite set $V \subseteq A^{Q \cup P}$ such that

- (i) $(v_1 \otimes \dots \otimes v_k) \cdot \mu'_k(\sigma) \in \langle V \rangle$ for every $\sigma \in \Sigma_k$ and $v_1, \dots, v_k \in V$, and
- (ii) $v_1 F = v_2 G$ for every $(v_1, v_2) \in V$ such that $v_1 \in A^Q$ and $v_2 \in A^P$.

With such a finite set V we can now construct a wta $M' = (\Sigma, V, \nu', G')$ with $G'_v = v F'$ for every $v \in V$ and $\sum_{v \in V} \nu'_k(\sigma)_{v_1 \dots v_k, v} \cdot v = (v_1 \otimes \dots \otimes v_k) \cdot \mu'_k(\sigma)$ for every $\sigma \in \Sigma_k$ and $v_1, \dots, v_k \in V$. It remains to prove that M' simulates $M + N$. To this end, let $X = (v)_{v \in V}$ where each $v \in V$ is a row vector. Then for every $\sigma \in \Sigma_k$, $v_1, \dots, v_k \in V$, and $q \in Q \cup P$, we have

$$\begin{aligned} (\nu'_k(\sigma) X)_{v_1 \dots v_k, q} &= \sum_{v \in V} \nu'_k(\sigma)_{v_1 \dots v_k, v} \cdot v_q = \left(\sum_{v \in V} \nu'_k(\sigma)_{v_1 \dots v_k, v} \cdot v \right)_q \\ &= ((v_1 \otimes \dots \otimes v_k) \cdot \mu'_k(\sigma))_q = \sum_{q_1, \dots, q_k \in Q \cup P} (v_1)_{q_1} \cdot \dots \cdot (v_k)_{q_k} \cdot \mu'_k(\sigma)_{q_1 \dots q_k, q} \\ &= (X^{k, \otimes} \cdot \mu'_k(\sigma))_{v_1 \dots v_k, q} . \end{aligned}$$

Moreover, if we let X_1 and X_2 be the restrictions of X to the entries of Q and P , respectively, then we have $\nu'_k(\sigma) X_1 = X_1^{k, \otimes} \cdot \mu_k(\sigma)$ and $\nu'_k(\sigma) X_2 = X_2^{k, \otimes} \cdot \nu_k(\sigma)$. In addition, $G'_v = v F' = \sum_{q \in Q \cup P} v_q F'_q = (X F')_v$ for every $v \in V$, which proves that $M' \rightarrow_X (M + N)$. Since $v_1 F = v_2 G$ for every $(v_1, v_2) \in V$, we

have $G'_{(v_1, v_2)} = (v_1, v_2)F' = v_1F + v_2G = (1+1)v_1F = (1+1)v_2G$. Now, let $G''_{(v_1, v_2)} = v_1F = v_2G$ for every $(v_1, v_2) \in V$. Then

$$(X_2G)_v = \sum_{p \in P} v_p G_p = v_2G = G''_v = v_1F = \sum_{q \in Q} v_q F_q = (X_1F)_v$$

for every $v = (v_1, v_2) \in V$. Consequently, $M'' \rightarrow_{X_1} M$ and $M'' \rightarrow_{X_2} N$, where $M'' = (\Sigma, V, \nu', G'')$. This proves the next theorem.

Theorem 9. *Let M and N be equivalent. If there exists a finite set $V \subseteq A^{Q \cup P}$ with properties (i) and (ii), then a finite chain of simulations joins M and N . In fact, there exists a single wta that simulates both M and N .*

Let us first recall a known result [2] for fields. Note that, in comparison to our results, the single wta can be chosen to be a minimal wta.

Theorem 10 (see [2, p. 453]). *Every two equivalent trim wta M and N over a field \mathcal{A} can be joined by a finite chain of simulations. Moreover, there exists a minimal wta that simulates both M and N .*

We can obtain a similar theorem with the help of Theorem 9 as follows. Let \mathcal{A} be a NOETHERIAN semiring. Let $V_0 = \{\mu'_0(\alpha) \mid \alpha \in \Sigma_0\}$ and

$$V_{i+1} = V_i \cup (\{(v_1 \otimes \cdots \otimes v_k) \cdot \mu'_k(\sigma) \mid \sigma \in \Sigma_k, v_1, \dots, v_k \in V_i\} \setminus \langle V_i \rangle)$$

for every $i \in \mathbb{N}$. Then $\{0\} \subseteq \langle V_0 \rangle \subseteq \langle V_1 \rangle \subseteq \cdots \subseteq \langle V_k \rangle \subseteq \cdots$ is stationary after finitely many steps because \mathcal{A} is NOETHERIAN. Thus, let $V = V_k$ for some $k \in \mathbb{N}$ such that $\langle V_k \rangle = \langle V_{k+1} \rangle$. Clearly, V is finite and has property (i). Trivially, $V \subseteq \{h_{\mu}(t) \mid t \in T_{\Sigma}\}$, so let $v \in V$ be such that $v = \sum_{i \in I} (h_{\mu}(t_i), h_{\nu}(t_i))$ for some finite index set I and $t_i \in T_{\Sigma}$ for every $i \in I$. Then

$$\left(\sum_{i \in I} h_{\mu}(t_i) \right) F = \sum_{i \in I} (\|M\|, t_i) = \sum_{i \in I} (\|N\|, t_i) = \left(\sum_{i \in I} h_{\nu}(t_i) \right) G$$

because $\|M\| = \|N\|$, which proves property (ii).

In fact, since $M + N$ uses only finitely many semiring coefficients, it is sufficient that every finitely generated subsemiring of \mathcal{A} is contained in a NOETHERIAN subsemiring of \mathcal{A} . Then the following theorem follows from Theorem 9.

Theorem 11. *Let \mathcal{A} be such that every finitely generated subsemiring is contained in a NOETHERIAN subsemiring of \mathcal{A} . For all equivalent wta M and N over \mathcal{A} , there exists a finite chain of simulations that join M and N . In fact, there exists a single wta that simulates both M and N .*

Note that \mathbb{Z} is a NOETHERIAN ring. More generally, every finitely generated commutative ring is NOETHERIAN [23, Cor. IV.2.4 & Prop. X.1.4].

Corollary 12 (of Theorem 11). *For all equivalent wta M and N over a commutative ring \mathcal{A} , there exists a finite chain of simulations that join M and N . In fact, there exists a single wta that simulates both M and N .*

Finally, let $\mathcal{A} = \mathbb{N}$ be the semiring of natural numbers. We compute the finite set $V \subseteq \mathbb{N}^{Q \cup P}$ as follows:

1. Let $V_0 = \{\mu'_0(\alpha) \mid \alpha \in \Sigma_0\}$ and $i = 0$.
2. For every $v, v' \in V_i$ such that $v \leq v'$, replace v' by $v' - v$.
3. Set $V_{i+1} = V_i \cup (\{(v_1 \otimes \cdots \otimes v_k) \cdot \mu'_k(\sigma) \mid \sigma \in \Sigma_k, v_1, \dots, v_k \in V_i\} \setminus \langle V_i \rangle)$.
4. Until $V_{i+1} = V_i$, increase i and repeat step 2.

Clearly, this algorithm terminates since every vector can only be replaced by a smaller vector in step 2 and step 3 only adds a finite number of vectors, which after the reduction in step 2 are pairwise incomparable. Moreover, property (i) trivially holds because at termination $V_{i+1} = V_i$ after step 3. Consequently, we only need to prove property (ii). To this end, we first prove that $V \subseteq \langle \{h_{\mu'}(t) \mid t \in T_\Sigma\} \rangle_{+, -}$. This is trivially true after step 1 because $\mu'_0(\alpha) = h_{\mu'}(\alpha)$ for every $\alpha \in \Sigma_0$. Clearly, the property is preserved in steps 2 and 3. Finally, property (ii) can now be proved as follows. Let $v \in V$ be such that $v = \sum_{i \in I_1} (h_\mu(t_i), h_\nu(t_i)) - \sum_{i \in I_2} (h_\mu(t_i), h_\nu(t_i))$ for some finite index sets I_1 and I_2 and $t_i \in T_\Sigma$ for every $i \in I_1 \cup I_2$. Then by $\|M\| = \|N\|$ we obtain

$$\begin{aligned} & \left(\sum_{i \in I_1} h_\mu(t_i) - \sum_{i \in I_2} h_\mu(t_i) \right) F = \sum_{i \in I_1} h_\mu(t_i) F - \sum_{i \in I_2} h_\mu(t_i) F \\ & = \sum_{i \in I_1} (\|M\|, t_i) - \sum_{i \in I_2} (\|M\|, t_i) = \sum_{i \in I_1} (\|N\|, t_i) - \sum_{i \in I_2} (\|N\|, t_i) \\ & = \sum_{i \in I_1} h_\nu(t_i) G - \sum_{i \in I_2} h_\nu(t_i) G = \left(\sum_{i \in I_1} h_\nu(t_i) - \sum_{i \in I_2} h_\nu(t_i) \right) G . \end{aligned}$$

Corollary 13 (of Theorem 9). *For all equivalent wta M and N over \mathbb{N} , there exists a finite chain of simulations that join M and N . In fact, there exists a single wta that simulates both M and N .*

For all finitely and effectively presented semirings, Theorems 10 and 11 and Corollaries 12 and 13 also yield decidability of equivalence for M and N . Essentially, we run the trivial semi-decidability test for inequality and a search for the wta that simulates both M and N in parallel. We know that either test will eventually return, thus deciding whether M and N are equivalent. Conversely, if equivalence is undecidable, then simulation cannot capture equivalence [16].

References

1. Abdulla, P.A., Jonsson, B., Mahata, P., d'Orso, J.: Regular tree model checking. In: CAV. LNCS, vol. 2404, pp. 555–568. Springer (2002)
2. Alexandrakis, A., Bozapalidis, S.: Représentations matricielles des séries d'arbre reconnaissables. Informatique Théorique et Applications 23(4), 449–459 (1989)
3. Béal, M.P., Lombardy, S., Sakarovitch, J.: On the equivalence of \mathbb{Z} -automata. In: ICALP. LNCS, vol. 3580, pp. 397–409. Springer (2005)

4. Béal, M.P., Lombardy, S., Sakarovitch, J.: Conjugacy and equivalence of weighted automata and functional transducers. In: CSR. LNCS, vol. 3967, pp. 58–69. Springer (2006)
5. Berstel, J., Reutenauer, C.: Rational Series and Their Languages, EATCS Monographs on Theoret. Comput. Sci., vol. 12. Springer (1984)
6. Bloom, S.L., Ésik, Z.: Iteration theories: The Equational Logic of Iterative Processes. Springer (1993)
7. Bloom, S.L., Ésik, Z.: An extension theorem with an application to formal tree series. *J. Autom. Lang. Combin.* 8(2), 145–185 (2003)
8. Bozapalidis, S.: Effective construction of the syntactic algebra of a recognizable series on trees. *Acta Inform.* 28(4), 351–363 (1991)
9. Buchholz, P.: Bisimulation relations for weighted automata. *Theoret. Comput. Sci.* 393(1–3), 109–123 (2008)
10. Cleophas, L.: Forest FIRE and FIRE wood: Tools for tree automata and tree algorithms. In: FSMNLP. pp. 191–198 (2008)
11. Eilenberg, S.: Automata, Languages, and Machines. Academic Press (1974)
12. Ésik, Z.: Axiomatizing the equational theory of regular tree languages. In: STACS. LNCS, vol. 1373, pp. 455–465. Springer (1998)
13. Ésik, Z.: Axiomatizing the equational theory of regular tree languages. *J. Log. Algebr. Program.* 79(2), 189–213 (2010)
14. Ésik, Z.: Fixed point theory. In: Handbook of Weighted Automata, chap. 2, pp. 29–66. EATCS Monographs on Theoret. Comput. Sci., Springer (2010)
15. Ésik, Z., Kuich, W.: A generation of KOZEN’s axiomatization of the equational theory of the regular sets. In: Words, Semigroups, and Transductions. pp. 99–114. World Scientific (2001)
16. Ésik, Z., Maletti, A.: Simulation vs. equivalence. In: FCS. pp. 119–122. CSREA Press (2010), preprint available at <http://arxiv.org/abs/1004.2426>
17. Hebisch, U., Weinert, H.J.: Semirings—Algebraic Theory and Applications in Computer Science. World Scientific (1998)
18. Högberg, J., Maletti, A., May, J.: Bisimulation minimisation for weighted tree automata. In: DLT. LNCS, vol. 4588, pp. 229–241. Springer (2007)
19. Karner, G.: Continuous monoids and semirings. *Theoret. Comput. Sci.* 318(3), 355–372 (2004)
20. Klarlund, N., Møller, A.: MONA Version 1.4 User Manual (2001)
21. Knight, K., Graehl, J.: An overview of probabilistic tree transducers for natural language processing. In: CICLing. LNCS, vol. 3406, pp. 1–24. Springer (2005)
22. Kozen, D.: A completeness theorem for KLEENE algebras and the algebra of regular events. *Inform. and Comput.* 110(2), 366–390 (1994)
23. Lang, S.: Algebra. Addison Wesley, 2nd edn. (1984)
24. May, J., Knight, K.: TIBURON: A weighted tree automata toolkit. In: CIAA. LNCS, vol. 4094, pp. 102–113. Springer (2006)
25. Milner, R.: A Calculus of Communicating Systems. Springer (1980)
26. Park, D.M.R.: Concurrency and automata on infinite sequences. In: GI-TCS. LNCS, vol. 104, pp. 167–183. Springer (1981)